RECORD

6

Collection and Field Note Ident

No. 15

(June 13, 1346 - July 11, 1946)

(asees subus)

		BOULDS! EXECUTES
		to the state of th
¥.	EHS	
	PS-5	** ACCORDANCE FRANCE CONTRACTOR OF LIVE DESCRIPTION
-	1E-85	
		The same of the sa
	NE-DE	The state of the s
	35-4W	ANALYSIS IN THE STATE OF THE ST
	PAUL PR	
	55-61	Wear airetrin lote Interded
	3-09	A TO THE PARTY OF
	15-60	White the same to the s
		Marigan, Manualton (Valley of Jetal.
		17 000
	76-79	here to I want the a management
	28-8S	He Description was a second of the second of
	34-35	OFFICE OF DESIGNATION OF THE PROPERTY OF THE P
	150-88	
		The same of the sa
	106-106	The state of the s
	100-SR	Annual Principle of the Color o
	200-10	
	101-88	
		A STATE OF THE PARTIES OF THE PARTIE
		Sesso-Alexan de learner les anno a Sesson de Sesson de les anno de
	16-301	Syste-Mars Smalel meglad to mordql-morel
	1-1105	Address of the contract of the
	J. Cak	Course of the state of the stat
	12001	The same of the sa
	Sin () 7.	ISEPS-0056S was asset to stude
		boo giff measure unestable, you accept to dived
	15-31	
	NEWS.	
		The same of the sa
	28+14	I william in the second

Collection and Field Note Book

No. 15

(June 19, 1946 - July 11, 1946)

(24942 -25526)

South of Agfayan Bay, plateau between Ylig and

Valley, Pago Bay 25355-25358

MARIANAS ISLANDS Page No.	MARIANAS ISLANDS	Page No.
Rota Island Airport	Guam, Merizo, south of Talafofo Bay, Inarajan, Umatag 25359-25426 Camarianas, hills north of Mt. Tenjo and Mt. Alutom 25427-25438 Description of Guam	130 - 153 152 - 155 154 - 162
Saipan, Kannat e Edtot (Valley of Ants)	to 25497), outside of village	166 - 173 174 - 185
Mt. Tapotchan, Saipan Island 25176-25198 78-83 Mt. Petosukara, Saipan Island 25199-25214 84-87 Garapan, Mt. Marpi, Saipan Island 25215-25225 88-91 Near Kobler field, Saipan Island 25226-25229 90-93 Mt. Tapotchan 25230-25232 92-93	Plants collected by Karl J. Pelzer on Truk May-August 1946	188-195
Agingan Point, Saipan Island 25233-25257 94-99 Tanapag, Saipan Island 25258-25263 98-101 Charankanoa, Lake Susupe, Saipan Island 25264-25276 102-105 Description of Saipan Island 106-109		

128-131



PIONEERS SINCE 1831

Miniature
Account Book
No. S 136

UNIT RULED

Made in 200 and 300 Pages

Made in U.S.A.

TO REORDER, SPECIFY NUMBER, RULING AND THICKNESS INDICATED ON BACKBONE OF THIS BOOK.

Collection number book of F.R. Fosberg # 15 to nos. 24942 - 25526 Descriptive paesages rerographed for file 5. Beefer

2 1946 - Marianas Is. Rota 3 native names and information from John through Juan Perez as interpreter - several others incl. Comissioner Ben in conference. June 19 - Rota I. airport limestone glat dense small tufte 24942 Eragnostis common, widespread on island 43 Ipomoea extensive prostrate blue, Istill ofen common covering ground, widespread on island "fofgo juice squeezed and used to at about 11 am. cure cracked feet, both human and cours, Tips crushed and applied to cuts. June 19 Lonson, Rota). persetting. firmestony soil. low isthmus, formerly site of town, destroyed Tripliasia tripolia forma Luing war, Many

plant, war, Many

acolypha willesiana etc.

44 Polanisia reo sandra demo dense thickete everywhere. herbogestrory seented, flowers -yellow. "mungus paluma" - bird food. seeds eaten by birds. common weed in old cultivated ground 45 Wealypha indieg L. styles white common weed in old cultivated ground Tuber "humatae" tuber 46 Eypens rotundus L. dense patch in old chewes for toothache. cultivated ground 47 quettarda speciosa L dump of slump "pano" wood used as sprouts arvind lange stump, flowers white. building material.

Roty 5 1946 Marianas 95. 24998 Pithesellobium dulce small trep 6 mm. Call ble greenish white, common everywhere fruit fendant on "kamachili" balls used for tanning; fruit eaten; leaves long pedundo. pines adged and used existed as emetic for dunk. enness, leaves chewed as remedy for two thache. large vine, older 49 Mucuna gigantea stems woody, common, climbing "badyogu ngadzi" vines used up to v cm, or more in diameter, grade pendent. medicinally to refresh insides and cut sections to supply drinking water. leaflers, stems 30 Cassytha fullformir C. yellow-green, flower orianional, parasitie "mai agai included in a sheumatism on busher of various kinds. 51 Leptures refrenz var. subulatua Fosb. loose, spreading ly numer. laraga finn rised as cattle feed. 52 Morinda citrifolial. small tree 4 m. Call flowers white, fruit common refrening unevenly lata wood used as building material; part of fruit sometimes noot for daying clother tan, fruit and white while affect is not used medicinally, mixed still flowering. with other plants in backache remedy. small tree 4 hm. tall; 53 Merserselmidia argentea leaves fleshig, silvery green; "burn" wood used be diving goggles, leaves will and weiter used as remely in case of prisoning from fish (tagoti, shippich) which have later point developed bruits fleshy.

Rota 1946 Marianas 14954 Clerodendrom low serambling show abundant, forming tangler between trees rup to 1.5 m. tall; Hervers white with fruglo stærnens. "lodugan leavesvery bitter, used for backache remedy. tree 5 m. Call; flower 55 albirria lebbels fregrant with slamens yellow common, possibly toward base, green "blackas" (because of moise it makes.

"blackas" (because of moise it makes.

wood used for construction. distally; pods rattling small tree 4 mm, tall, Il Melanolepis multiglandulosa pendent of fruits , common in thickets 'alum' young tips eaten; back used medicinally to refresh inside rounded should 1 mg. 57 Leaevola frutercens (Mill.) Kr. tall, flowers white. common near beach nanasa ripe fruit used for sone eyes. showb 3 m. tall, 38 Rophera tomentora occasional near boach fruite dry, brittle 39 Fimbristylis dense clumbs, leaves "tutopu sabana" noote used medicinally juice agreemed and used in buby; mouth for coated or cracked tongue. very common, these plants from shade LO Prema obtraction R. Br. (printer los) showb 2.5 m. Call, "ah gan" wood used for Givewood and construction, leaves medicinally for flowers green, aromatic. refreshing insides a releasing heat" should im talf. Ils. white geen early. 61 Prychotria mariannensis as building material, ripe fruit eater by bride

Rota 1946 Marianas S. 24962 nostor partially dried when 63 Barleria cristata
ran, persisting from cultivation collected, collapsed. should, m. Call, flowers o malex, coally readily falling. June 20, same, 64 Portulaca oleracea L. plant prostrate, common awind camp bleshy; flowers yellow, about , cm. in cultivated ground budu lagar fed to pigs across, ofpen in and boiled and eaten by mnning, petals leeply emarginate; humans. stamens 12-15. 65 amaranthur spinosust. common in disturbel places stem ascending "bilitis" young growth eaten by humans a fed to prays. 66 grewia small tree 5 m. tall, around develling flowers green, slightly fragrant. 67 Tragnostis amabilis L. spreading lifts common expecially on sandy disturbed soil. 68 Eughorbia hinta L. spreading assending, very common in abundanced greling at tips. "golondrina" used with "abas" or cultivated ground. Pridium quajava for thyllanthus

10 1946 Mananas 90. Rota 11 amarus Schom. . Th. "maigo lalo" used with 24969 Phyllanthus mini at. G. L. We botu, 1955 Englished hirta as vagend astringent common everywhere in also & relieve rectal itching disturbed places leaves mashed and inserted. saf milley; flowers 70 Hippobroma longiflar pure white. rare in abandoned cultivated ground. sheather strongly glattened. 71 Chloris radiata common everywhere in cultivated ground, rvadsider, etc. June vo - n.w. evast on second terrace the sol is how and very of mest end of main part of Rota Island thin. The Japanese raleed it tera benefit of coral up into windows to get depth enough for siveet limestone separated potatoes and Whanihot. by cliffs. 72 Fiers nearbase of eliff Type tree, fruit green. 73 Ximenia americana thickets on second Terrace low langled showl, 1.5 m. tall, fruits yellow when rife, fleshy, rather sorry not of especially good flavor. "pint" fruit enten bychildren when ripe.

20 1946 Marianas Jy. 15002 Hedyotismariannensis Rota sub-shrub, frelid when broken, fls. white; spar dedo" medicino for margina disease believed to be contracted in the jurgle and caused by anger of the sprits of the ancient Chamoreus characterized by pain, fever, swelling and delirium at night. shub up to 1.5 m. 03 & hyllanthus and clifts tall; gaagas wehan medicine for backacho. banyan-type tree, fruit gree minu used mixed occasional with vinegas for internal frains or & our outs. nouts used used as flirewood in line bilm. 0; Bilskig manannense cliffe, common showly write white flowers. gan sali" word ward for trebe and firewood. 06 thickets "anglao" (pronounced angilo) used for grage for cattle.

marianas)s. 24 1946 Rota 250/3 Piper woods in raviso on cliff should Im. tall "Though and to care genorleg. strub "sumag" forest in elift used medicinally, ete.

mall tree, "tatte 15 Pandanus tectorius common in forest "cafifo" or kafri Cave, used in bastactry, fruit and mute sates. Juis. "Batod" juice 16 in frest said to cause severe sken initation and rash, but said to be used to cure tous. show "sadafu" 17 weed in fields 18 Delringia amaranthads, frest on cliffs back used for cordage "pechalan" novy used in mixture "to refreshinsieles." low should, fruit red. "agatelong" wood 19 Engenia palumbis forest on second terrare very flexible mselfer axe bandles, frosts for bullerle cart, etc. fruit eaten.

Rota 26 1946 Marianas 75. 4. 'lalaha' wood in frest 25020 torsles, etc., buins readily without small tree "paipai" 21 quamin saffordis in firest or second terras word very flixible, used for handles for utensil, tongues for bull carts, building materials, etc. va Emphorbia heterophylla bracks red at base "flores hagin" latex initating to skin shrub 1-3 m. lale 23 keneaena glanca omme everywhere woods "talntangan" word used as finewort a bouilding material. seed used during way as coffeesulstitute but caused loss of hair, leaves good frig feed. Tree "derog" woord weed 24 Calofolylluminsphyllum papanen temple ground bull-cart where. and bouts. lust well in water.

32 1946 Marianas 95. 25038 Ardisia vecasional

39 asplenium epiplingtie on The trunts

40 asplenium epiphytic on tree trumbs

41 fregginetia

42 asplenium nidus very common, mostly epiphytic

43 Matteton Classylon common

44 Dioseorea bulbigera

45 Hernandia lahyinthica Tryanni common in forest

June 72 same
slopes of volcanic breccia
underlying limestone
cliffs. slopes grasscovered, badby eroded
in places, bath by
waters and slippage,
showing evidence of
burning over.
25087 Mytella benningserians

local on enoded place

- 78 fphenomenis eroded places
- 89 Frirena muddy seep
- 90 Colocasia esculenta planted in moist placer
- 91 Elephantopus seaber common along trace
- 92 Dryopteris goggilodus:

Eyporpordium ternum gleichenig linearis common on eroded places, Miseanthus japonicus dominant on slopes

dense bushy shrube up to 0.7 m. tall, fols. white, fruit reddish black when rips.

stifre, red, fronds

sentagonal.

bluish cast.

46 1946 Marianas Is. Rota Traniele drooping Onyza sativa in small muddy place above trail stiffly erect lights. 94 Fimbristylis vecasional heads with white 95 Brdens pilosa var mina rare along trains low shoul ; flowere white, 96 Scalvola frutercens (Milf.) Kr. oceasional fleshy. 97 Heterofrogen contestus rare flowers purple. 98 Emilia somelifolia sare along trail herburget in to 1 m. tall, flower yellow. Islander trifts 99 Cassia rare along trail 25100 Ryndrospora occasional in eroded places. 01 Eleveloris muddy seeps flowers yellow 02 Lynedrella modiflora disturbed place strictly erect, stip nodding, plowers white.

flower yellow. 03 Euphorbia 04 Curenligo eroded burned place common locally

52 1944 Marianar 90. 25122 Proces pedunculata epiphytic on tree trumba just back of beach 23 Cyclophorus common on brancher and trushes of trees just 24 Digitary minimum (Kunth) Ham common on welsy beach and base of cliffs 25 fleurya mderalis 16 Clerodendron great rock on base of clift 27 frest, common 28 - 150-100 frest, common 79 Hernandia ned comment trees on frest.

Rota 58 \$1946 Marianas 90. wordy ving, buds 25143 Alyxia spiphytic in large tree and green fruits only. rarely fruiting. Thee trunks.
45 Piptuus insanus
common in grest small tree, leaves dark green above; fruit very juicy, white, acheneron surface. tree 4 m. tall, leaves 46 Pandanus common in frest sprinally arranged, fruit not quite ripp. tree your tall bushy, 47 quamia saffordii common in frest llaver clarks green, fruits green, turning red, then black when ripe. herb 1 m. tall, flowers 48 Clerodendon speciosissimum scarlet, fruit immature persisting in tiny abandoned greenish black. clearing; japanese compraite. herb up to I m tall, 49 folanum nigrum Tiny abandoned cleaning flowers white, night fruit blue-black. servi- seandent shoul 50 alycia thry abandoned clearing flower dull salmon flower dull white. epiphytic on Tree truck, race

Rota 1946 Marianas Is. Tree 5 m, tall, fruit 25152 Pandanus 53 common in frest still green. fleshy, leaves roft, occasional in frest not stinging. 54 Heffetlefris Belvina spicata rare, epiphytic on tree trunks June 23 - commit grove beyond phosphate mill, on road to as malote 55 Parpalum spreading 56 Digitaria priviers var. pruviers 57 Paspalum conjugatum except along wasted plain June 24 - near the airstrife, near east end of Rota I. small pratch of woods in scrubby second-growth in terrace should 3 m. tall, young leaves whitish from publishers, very abundant in semb flowers day frink.

one where

Rola is a series of coral limestone terraces about five of them I the second from the top is a but confused when viewed from The north, with a low tip of volcanic tuff, projecting thru the middle of the topmost, and a considerable exposure of volcane breises in the southeast side, this capped by a thick layer of limestone. The third main terrace is the most extensive, and is wherethe air stry is situated. actually there may be several more minn terracer than the 5 mentioned, but they are hard to separate: The island has more vegetation than I iman, apparently because there is less arable soil. On exprosures of telling malaster the original frest fressists in fairly undraturbed

66 1946 Marianas Is.

M. wens, are tangled in the trees. I lagellaria ending and Frenzeinetia are emmon climbers. tagraea sp. is not common. It has a strangling habit

It is hard to tell if this frest was what was found on the parts of the rupper teraces whee soil was most abundant or not. These places have been

have been planted. Under these nothing grows. Most

abandoneg. Pastralum

and folanum nigrum

as well as Crotalaria

stelecantha (?), quamid -saffordii, Macaranga sp., Classylon (???) with frama,

andisig of. Piper sp.,

as shrubs. Mucuna gregantes and (more rurely)

as in frees and Clusia.

cleared. Wind breaks and

patches of acacia confusa

of the cleared land is now

conjugatum, V. orbiculare

are abundant rokeds,

(25158) las come in in almost pure stand over large areas, forming a servet 2-3 m. tall. I'm open places l'assiflara foetida or pornoea sp. (24943) a both from a dense mat overthe ground. On the lower terraces the strand character of the regetation becomes more and more pronounced. the portions of the second and its associated terrace examined had been largely cleared, except for patcher of serubly freit. (his that Inthis byings, Friend tenctorin, Kinenia americana, Prema Gymnospinia, Pandanin fragrans, milandehis P. Lexitrius, Guarnia Hibicus and others. The Japanes Ochrisa had planted Casuarina in palebre. Sweet polatoes, casava, and papayar are persisting from cultivation. On places where the soil is only a few very then

the Japanese had raked it uf in windrows following the contour and planted on there. The ciral bedrock as is exposed between these. the lowest tenace forms a sort of crastal plain along the northwest ovast, at least. It hes, here about 2 m. above sea level. on the southeast court it is much higher, 10:15 m. a more. On the isthmus it is variable. This upparently had, originally, a strand forest of Hernandia (sp. with peltate leaves), Thespesia, Hibiscus tiliaceus, Barringtonia asistica, l'andanus tectorius, landanus fragrans, Terminalia eatappa, Clasicylon (???), Laporteg (?), Macaranga, Fremna, Ochrosia, guetlanda speciosa, Pithecellobium dulce (an important component but probably introduced),

allophyllus, Piptrems argenten, morinda celrifolia Melanolepis multiglandulosa, occasional Pismia grandis, albiggia lebbets, grewig, oriasional Muntingia calabra, Fiers 2 sp., Intsia bijuga (common), along the feach Menerschmitis argentes, Sophora tomentora, Exroesaria agallocha and abundant Lesevola futerini undergrowth of Phylanthus Janninum, Clerodendun totulring asiating, Callicarpa, & and, above all dense thishets of Triphasis trifolia, the whole forest entangled in Mucuna gigantes and, more rarely, M. revens (or princene), with spornous alla near the orean. In opening are many weeds - thyllanthur ! mirini, healyphase. Englishing forty, Etrymifolis Cassiy occidentalis, sidy of (aff. nhombifolia), Elevin indieg, Ehleris radiata, Lotanum nigrum, Volanisia

Digitaria sanguinalis,

Cyperus notunctifolius

forming dense patches,

C. stuppers in isolated

tufts, common, Lepturus

repens near the coast,

Thuranea just at top of beach.

asslepias curassancus, Euplustic

heterophylla

an cliffer and big rocks near ocean Cemphis acidula conly right near ocean), Bildsig, Terminalia sp., Canthium sp., Clevodendron, Wedelia, Digitaria mariannensis, Davallia solida, asplenium 2p., Psilotum nudrem, Hedyotis albido-frunctata, and oit bases of clifts Hedryotis foetida van mariannensis and a fern (25138), a large Dryspteris (25137). On the branches of trees Hypmenolefis spirata, Davallia solida, and Proces pedunculata, and

a sterile Elaphoglossum grow epiphylically. On the brigher terrace on the south coast Lodonala viseosa in common also Parfealum enjugatum. Coronits have been common along most of blowder the coast, but except for one or two places only stumps remain. Where any trees semain they by a weeril. Cultivated frants of various sorts gresset on the lower terrace -Citrus limes, C. amantifolia acalypha willeriang agave sisalana, a. americana? Calofehayelum inophyllum, Rhoer discolor, Polyscias quilforglei, the Myllanthus nivosus, Codiacum variegatum, Rieines communis, Caries papayalone of the most abundant weeds), Bougainville, Delonix regia (common), etc.

on the south coast above the village of as malste, is an extensive stope of a deeply weathered volcanie breezig. The This is open country, frequently burned, and with large error and slippage sears. I has a distinctive flow. the dominant plant is Miseanthus japonicus. other things are chiefly found in and around a serving and west seeper al, Rynchospura sp., Friena of., Finbristylis of, Curculigo sp., Euphindia of., Heterofiogen contestus, Clerchain of. Notholaena sp., Aphenomeis sp., Tacca lemtofrodivides, and skirubs of Wilestroemia elliptica, genisstoma sp., and expecially # 25087. Tyredrella modiflora is a prominent weed in cultivated areas in this region. In a small marshy place was a

This slope

77 Laipan 76 1946 Marianas Is. aval limestone with June 27 - Kannat a Editot (Nalley of ants), 3 km. e. of afetina Pt., Saipan). small vally, abandoned coffee plantation, with Vassiflara foetrda smother. ing out the coffee trees, hee 5 m, tall, with 25164 Enythina small prickles. occasional tree 6 m, tall, leaflets 65 aglara treatending to droop; cliff, on talus. flowers crange, fruit range when ripe. small tree, fruit of Cynometra immature. small at growt of eliff 66 Melanolepis multiglandulos shrub v motall, builing spikes pendent emmen pendent should 68 alyxia corolla love yellow, common of on frace of tube crange small treds 3 mitell limeston eliff. 69 Coffee andring L. ripe fruit red. es plantation, formerly cultivated tree & m. tall, rounded, 70 acacia confusa bushy, flowers plantey on windbreak gellow, fragrant, thee 8 mitall, "flowers 71 Intria bijuga common in valley with green calyx, nings teaming and style farmises

Marianas 2. 78 Laipan V5172 Passiflorg foretida var. hispida 172 Killip very abundant everywhere herbaceous vino climbing over everything, Mange. Mange. 73 Leucaena glanca very abundant in for the shrub 1-2 m, tall. inthickets and invading ofpenings forming a deep soft, long 74 Eynodon daitylen va. mat about 3dm, deaf. bery abundant in opening small tree 5 m. tall, 75 albyzia lebbels meading, glowers common everywhere cream yellow This specimen in raving June 77 - south ridges of mt. Tapotchan Chighest Mountain on island), faifran. rough limestone knot and slopes below it. thickets and openings should 15 mitall, flowers 76 Psychoting mariannensis white, ripe fruit occasional on brushy slope ald. (& plant is well.) show who to Im. tell 77 Phyllanthus flowers greenish white. very common on nocky slopes and bluffer. showly you tall frint 78 Classylon(2) green, fleshing. occasing in thickets leulms solitary, 79 Cypeus sprhelets yellow your common of fren alopes

83 Lailan. 82 1946 marianas 95. flowers yellow. 25187 Curentigo common along old road caespitore beaditan. 78 Fimbristylis in common in enoded spot Tufts 89 Ryndrospors common in ended spot weal stemmed go feter pogen contatus common is evoded sport unewhat easysitory outmos erect. 91 Fimbristylis common in evoded and 97 letaria emman along ward rays white. 93 Bidens pilosa var minor small patch along road. spreading to prostrate 94 Edinochloa colonin disk yellow, raylers. occasional along road 95 Bidens pilos oure along was fls. yellow, standarp outside. 96 Claschynomene Sestania connalina occasional along road 97 Puera in Calopogonium muumoides Desv.
nave along road
98 Cassia
rare along road flowers yellow.

84 marianas Is. 1946 June 28 - sea cliffs east of mt. Petroubara, m. and cliffs and steep stopes mostly of a triff that is laigely composed of light gray pumice in a white coumbly making 25199 Wedelia liftere van. combiens bushon limestone alope 25 200 Scaerola frutescens (Mill.) Ken 01 Bikkia & ledger on tuff cliffs 02 Heteropogen contentus dominant in sports where Miscanthus is absent 03 fimbristylis cymora common in open places incliffes by Vicia of mat at top of limestone cliffe

87 Laipan - Rota 1946 Marianas 95. 86 rays yellow, short. 25205 Bidens glorogyne termifolia 06 Ipomorea alla L. prostrate, extensive, flowers white, closed roadside, large palely at 10:30 a.m. twiner, leaves strongly 17 Cantharospermum scarabaeoide (L.) Baill reticulate, gray beneatly, climbing on shrubs on flowers dill yellow. open tuff stopes (seen also but sterile with \$ 25183-25198) clembing on buster, 08 alyxia ledges on tuff cliffs flower lemm yellow with wange tube, fragrant prostrate, flesly, og Portulaca gradrifida L edges of tuff cliffs flower yellow, about 15 mm. across. 10 zvysia snatrella postate erect lifts spreading by prostrate ofren tuff slopes and ledges prostrate, pleshy, II Hedystis albidopunetata crevises and ledges in tuff cliffs flowers white. Rota June 24 - same as 75110-25114. cultivated around in sorryard prostrate, purplish. 12 Eugherbia Migmefolia 13 Kilea merphylla densely covering ground causes mormally prostrate · parasitie on Cupholia thy mipoling

1946 Marianas 95. faipen 96 show hanging over Viter negundo common conal rock edge of roots, blower lavender. 41 Messerschmidig argentea sand dunes, common shrub 2.5 m. tall, leaves fleshy silvery green; flowers white. 42 Wedelig biflora van. camescens sand dunes prostrate, transles ascending, flowers Keaflers vine, green 13 Cassytha filiformis Howen white, saind duner, elimbing over and parasitic on quit green various strubs prostrate grining love 44 Thuarea involuta top of beach worly caesfritors 45 Cenclines echinature common on dunes and 46 Hernandtia. small tree 4 m. tall, leaves glossy, florous while. common tree along top of beach 47 Tacca leontopodivides small colony to in shelter of sand dune prostrate, palegreen. 48 Enphorbis sand dunes berb 0.7 m. tall 49 amaranthus spinosush. sand dunes glowere purple 50 Vernonia cinera sand dunes, common

98 1946 Marianas 91. Laipan prostrute extensively 2525: Spronoea per-caprae beach flowers purple closed in cefternoon 52 Walvastrum coromandelianum common on dunerand flats erect herb, brusts 53 Emphorbia heterophylla searled at base. bushy best up to 0.8 m. 54 Lida common on duner and flats tall, flowers dul range-yellow, falle toward outside. I specimen from several glants) prostrate, flowere 55 /riumfetta procumbens deep yellow. occasional in upper beach stems ascending, 56 Euphrbig hirta arching at tips. sommon everywhere tree 10 m tall, 25 cm. thick 57 Casuarina equisetifolia these trees looks quite common along replex healthy as do most edge of beach that and down along crast, as contrasted with those up in mountains) derostichum aueum June 29. Janapag, faipand. clumps among small mangrieve swamp partially filled in with with, mangroves. only one species of mangrove seen, other associated plants around edges,

Saipan 101 1946 Mananas Is. 100 tree 5 m. tall with 25258 Bruginera Dolunt, to vee - shaped dominant plant in colcareous frommatophnes; tidal med flat calyx bright red, glosey, yetals still glesh-nange; radical 17 & arrially binne noot) begaginal in cross section. white, fruits bringetel, 59 Hernandia mangrove swamp pale green, not quite clusters below flowering clusters at flowering time aifice circular, for plugged by from when immating. small tree 4 mitale, 60 Thespesia populnea flowers pale yellow common around edges center. deep fruple black of mangrove swamp vine, flowers purple, 6) pomoce small (rem. across) 'along wadside prostrate, extensive, 62 Vigna tate marina freduncles erect, along roadiides and rubble heaps flowers dull yellow. 63 Hibiscus Trliaceus edge of mangrove wasp

103 1944 Marianas Jr. Kaipan 102 Charankanva, frine og - west of Salve fumpe, tarpan. low ground in wastel plain should 1.5 m. tall. 25264 frence vow rounded spreading 65 fithecellobium dulce tree & milall with fence ow, commonent fruit pendent, coiled. lowland tree. tree 6 mitall, flowers 66 Terminalia catappa white, fruit very young. rvadside June 29 Latre Luxufre. faifran /. lake of clear dark word water, almost fresh, surrounded by marsher of Parpalum valginatum principally acrostichum arream, the odendron of. Hibiscus Tiliaceus ele. with a few large lamaring 67 Paspalum vaginatum pure stand forming dense truly on wet fround large clumps fronde 68 acrostichum auseuni common in swampy ground

faipan 104 1946 Marianas Is. 105 flower purple 15267 Sporney 70 Cleroderdun turf forming large masses. abundant in swamp from 29 - marshes and pends Lake Surupe, Laipan 7. 71 Justices of prind abundant in water I m. long, flowere yellow, petale obradate. caespitore 72 Cyperus culms terete 73 Leipus abundant in water shrub 3 m tall. 74 Indigofera suffritiona mill. occasional near prond. decumbent with ascending tiger, stems fistulose, plants sterile. 75 Hornoca reptans abundant in mud June og - afetra (fusufe) Pl. doorgand faifan). 76 Heliotropium indicum d. flo. pinkish white.

Jo faipan 107 manaras 1s. 106 1946 the island of Saipan is much less simple in its topography than Rota or Timan. Like Them it is principally of raised coral limestone, and a number of Thraces are in places evident. There has been much errsion and probably other disturber ance, resulting in a rather high mountain mass in the center (Mt. Tapotchau), continuing rather niggedly to the northandbut sloping off rapidly to a low ridge in the south. Much of the western part, except in the month, is a broad Crastal plain, variable in beight but not more than y m. a lake of almost fresh water and a string of grands and marshes lies aling the inner edge of foetida almost smothers everythingout. this coastal plain. around the late is an extensive swamp

,097 -Agricely An Lite ale J Block proporty) F1 -12+61 Partings From Mark AM

107 faifran marianas 1s. 106 1946 of acrostichum, Clerodendron the reland of Sarpan and Varfallum vagenation. is much less simple The marshes are filled with Rota or Timan Cily Leirpur, Jussacea, Homora regitans, Them it is principally This island is so a raised coral limestone, disturbed and torn up and a number of that it is little more Thraces are in places than or great weed patch. evident. There has been all of the lowland much errsion and has been under cultivation probably other disturber or bulldozed eneght ance, resulting in a the extreme strand, and rather high mountain even-this is not undisturbed mass in the center in many places. Most (Mt. Tapotchan), continuing of the ordinary tropical rather niggedly to the Iveeds characterize the northandbut sloping off lowlands, and are about rapidly to a low ridge Br. the only thing there. along in the south. Much of the beach and occasionally the western part, except inland are large Casuarina in the month, is a broad trees appearing quite Crastal plain, variable beatthy. Inphasia in beight but not more than trifolis curses the visland, y m. a lake of almost fresh forming thickets both water and a string of in the low and middle ponds and marshes ties altitudes. Passifling Har. aling the inner edge of fretida almost smothers this coastal plain. around everythingout. The large is an extensive swamp

109 Laifran 1946 Marianas Is. 108 and at their bases. Here are Conberg britispeloting, One large ridge to the south of my Tapotchan is Randia, Ficusto Lapentea, covered by Miseanthus and Pandenus (2 sp.), l'aptures, etc. 108 has a slight development of the On the bigher slopes of volcanic soil flora. The soil is the main mountain man thick and clay like, but underlain are also some fairly good by limestone. Otherwise the slands of native vegemiddle altitudes are tation, both on rocky either cleared, or thickets outcrops and slopes. of Imphasia, albezzia and Phyllanthus, Bibling, acasia confusa. l'atcher Rospelistria, and a number of remnants of native of trees are here. vegetation such as landanus Intsig bigingg is (2 sp.) and Freus sp. remain scattered almost everywhere here and there, but most as is Cynometra, but the of the brush and woods latter only on rocky places. are acacia and albyzia, alyxia is also common where which are apparently a trace of native vegetation persix competing for complete Volcania material comes prossession of the island. to the surface here and The Cleasia, expresially, forms therein the mountains, but a dense stand, permitting seems mostly covered nothing to live beneath it. Cy Parpalum conjugatum. Ce From a distance the mountain tiff exposure, gray pumice breise, appear wooded, but most with north-east side, has a of, it is there two species. 33 por development of the volcaine on the cliffs many soil flux, mostly Heteropogen native species persent. and Miseanthus. The boundary between frest and grass is very sharp on the centact . both actually in them

July 1, Mt. aluton, quam eroded slopes of volcanic soil, generally covered by Miseanthus japonieus, but enoded sports and shettered places have a perulia flua, simila but with more numerous species than that on Rota on volcame soil. feveral sedges, Leavola fruteriens Withstroemia, Dianella sp, Pityrogramma, Blechnum, Iteria, glochistion, Bidens

15177 Leaevola fruteriene common

78 Mitracaypum common

79 Eughorbia common

abundant

81 Desmodium triflorum ammer

82 Bidens glonogyne tennifolia extrared local, in crevices in Ged well

quam Euphorbia sto., Lycopodium comming glercheng linearis, etc. all characterize it, besides a number of weeds, as Emilia sugittata, Stachytagheta, to etc.

low should I m. tall, rounded; flowers white - buit white, fleshy. erect, flowers white. grayish green. prostrate, flower purple. " blowers yellow.

Gam. 117 116 1942 marianas Is should mits faither April Marker. black raises spots 25305 lichen acc. W. W. Dichl, sent to Claus. on barls of dying branch of aglaig (25303) small tree, erect, 06 Clarkeylon pale green. occasional in frest small shrub, ref \$ 0.8 (07 Hedyolis feetidus vas. m. tall, flowers white. 208 common on differ and plant foetid when broken. should 15m. tall, flo white. 10 Billing mariannensis vas. 11 Morinda gumbellatav. glandula woody ving, fruits -yellow when nearly ripe. climbing in trees in frest prostrate, mat- former 12 thrace involuta pentale common on level places 13 Elatostemma on inner face of rimof fronds erect 14 Polypodium seolopendrium edge of cliff or rough limeton tree 5 m. tall; flowers brilliant red orange, 15 Cordia subcirclata Cam inner face of rim of cliff bruit green. small-shrule, flowers 16 Billhia manannensis white. just below edge of cliff,

119 quam, 118 1946 Marianas S. showly 2 m. tall i flowers frink, deeper at center. 25317 occasional in edge of cleaning meaning cliff spreading tubte. 18 Engrostin 19 Phyllanthus common in well crevices 20 Engenia Cliff compact bush o. 6 mitall, rife fruit red with on rum of diff green calys. shrub, flowers white 21 Biskling (photo) just below rim of cliff July 2 - just south of Agfayan Bay, south of Inarajan, quam. reef flat, slightly exposed at low tide sterile, rhizaner buried in rand. sterile, stigemer deels in sund, leaves distichous, abundant all over rey flat mostly franced off at tips, Two lateral filters persisting after leaves not away

Mananas Is. grunn 1946 124 "marigrig" sometimes used medicinally." 25338 Wedelia treflera var. carescero with ascending handes common on sand shoul 2.5' m. talk 39 Leaevola friteriens fleshy "manaso" to prepare a lotion. emmon shoul & m. tall, leaves 40 Messerschmidig argentea occasional or sand silvery green flesly; flower white "unig" ka "lotea caespitoza leaver rentariona. 41 Fimbristylis
accasional on sand
42 Vigna maring
abundant on sand somewhat stiffish. prostrate, extension. pedundes erect flower naplor"
yellow. wedicinal formany ailments. semi- seandent should 43 Colubring asiatica feaver glorsy; flowere yellow. "gásoso" used medicinally for many ailments. common in thickets prostrate, stems green; 44 Lesuvium portulacastium leaves thicks fleshy; "chara" enter cooked as a vegetable. Hower finds within. leaver white beneath 45 Eupherbia Chaminonis Bois. flowers greenish. "betulagas unci" rare on sand

gream 129

small tree 4 m. Talp,

fruits green. "myolon"

wood used for house foundations."

semi-reandent shoul,

fruit reddish grees.

"laluput" used medicinally.

geory; flowers white growing, stigma horse-shoe shaped; fruiting perigone red inflated, when ripe, with round origine. Items ascending; leaves thin; flowers white, corolla tube to medicinally.

Low shoutly tree 4 m. tell; leaverdark green above; flowers pale yellow with maromants.

133 guam 1944 / Mananas)s. small tree 5 m. tall, 25361 Thespesia populnea flower pale yellow, "banalo" or "kiluli" wood used by house toundations; fruit used by children to make tops. back boiled down and was by women for small Tree 3 m. Call 62 Khizophora munonata 2 am with arshed profronts, occasional in mud Tree 8 m. tall, branchets drooping, 63 Casuaring equisetifolia gago' very hard wood used · common along beach for foundations for houses, also July 3 - just south of good for chancoal. leaves used for Talafofo Bay, quam brushy slope tangles amudging morquitoes at night, with wines, trush mostly Triphasia. were tangled over 63a Divseney medicine for fatigue and stomach ache. "gado" note laten; leaves used for small tree 5 m. Call; 64 Cording subradata Ram fruit immature. occasional wood used for house foundations small tree 4 m. tal 65 Premna obtusfolia L.G. (printed los. leave, dull, much occasional affected by a gall, flower, green "again" valuable word, as good as "fil" termiter don't like it. leaves used in medicinally. Endisease like preumonia. Leaevola frutereens July 4 Inanajan, s.o. wast also prominent. of quan disserted rough coral on edge of sea, mostly covered by Pemphis acidula,

136 1944 Marianes 25. quan showly am tall, flowers 75371 Jana white. cultivated as an arramental vine, extensively dint. 72 Monndiea charantia ing, plowers yellow, weed around building ofsen in morning, closed and thickets and dropping in afternoon; fruits redcrange, clastically dehiseed at touch, said to have been brought to againa from Priesto Rico; then the local priest, Father maring brought it to merizo before the war, planted it around the church. now it has spread ent the surrounding spreading tree, later thin abilidant, it yellow flowers green. Truits yellow 73 Jatropha cureas inthickets when mature, dry nute still banging on tree. "tuba tuba" safe used for Olowing bubbles; mits eater but only in a mall quantities, also strung and burned in place of carlles. "chaquan Rabajo 74 Eleusine indica L. freder for cattle. comma everywhere

139 quan 138 1946 Marianas 75. fleshy, depressed to ascending, 25375 Portulaca oberacea disturbed ground flowers yellow, stamens about 10. "bedulagas dangenlo eaten corbeel ær a vegetable. show i'm, tall, flowers 76 Cassia yellow, pods terete. probably planted, near dwellings. "amot tumaga" leaves used medicinally. flowers purple. 77 Vennong cinerea chaquan Santa Maria"! common weed "laha" for foodder 78 Paspalum occasional in open andring, "golondrina" 79 Euphobia hirta both leaves and roots very common used medicinally to euro cuts. 80 Phyllanthus sinus sohom. 2Th. Welste, 1955 erect herb "maigo lalo" used medicinally, plant common everywhere in open Goiled in water and water drunk for dysentery. 31 foldnisia ico zandra common in disturbed ground. bushy herb, flowers yellow. "mongoz paluma". used medicinally; seeds eaten by prigeons. vine clembing into lower 82 Diosessea cultivated branches of trees. "mika" tubers eaten. used medicinally for stomachache.

141 quam Marianas Is. 140 1946 25383 Tacca lemtopodioides " gab gab tubers used for making starch for cultivated (occurs both food for invalids planted and wild) (plants and "hamany" plant soabed 84 Hibiseus " in water and a muciloccasional aginous extract drunk for cooling effect by people in working in bot weather. also medicinal. vine, sterile. "hikamas" 85 Pachyshizus erosus tubers eaten.
vine; "dago agaga" cultivated. 86 Dioseorea aluta (red yam). Tuber eaten, cultivated scrafted and made into pastry. " vine "therile; dags haya" 87 Dirrerea alate cultivated (pronounced haija), tuber eaten, me of best yams, steile vine "dags apaca" 83 Da Dioserea alata 39 cultivated root eaten. extensive vine 90 Urena lobata
common weed in waste places "alamli" usedas fødder. "dadangsi agaga leave, and rooks brilled and used medicinally. brigame. "Kahalas" 91 Polybodium scolopendrium used medicinally.

142 1946 marianas Is. quam small tree "hoda used 25392 Freuz tineteria medicinally to treat children with coated tongue and throat as in diphtheria. of they blands well-framed tree 94 Calophyllum inophyllum "davg" or "pals maria" seattered wood used for bull-cart wheel hubs and axles. one of toughest woods available. Often used for carving. petioler pale green 95 Colocasia esculenta suni pasencia o cultivates in low place white taro; leaves and worked and eaten. semi-seandent shrub, 94 Clerodendon "lodigan boiled and inthickets water drunk to remedy fever. small dense tree, 97 Theretia permiana flowers bright yellow. or namental, semi-wild. sectives dell red purple. 98 Colocasia escutenta "suni agaga" or red two cultivated in low places mature leaves and sterile clumps "mumutung polo palavan used medicinally (distinct from 25406,

quam 145 144 1946 Marianas Ds. small tree with 25400 avershoa carambola L.
growing on edge of field, probably
planted red influences of vose purple flowers with white edges fruit with 5 strongrifs. belembines fruit eaten. large dense tree éarabas Of Mangifers indica scattered in field mango" best of local mangrer; fruit large, not fibrour. faid to be propagated by seed and to come true. amall tree, flowers or Morinda citrifolia L. very emmon white; lada leaves applied to the frehead to cure headache. vino, sterilo. "gaye" oz Mucuna gigantea common shoul, sterile " ufa" 04 Heritiera littoralis rare in low thricket hubs. erect, fleshy, sterile. 05 Brysphyllum calycinum -common along roadside "semperviva medicinal. erect clumps, somewhat lake" not used Chistinet from # 25399,

147 quam Marianas) s. 146 1946 deep tangles of very Triphasia trifolia 25407 spring bushes, fruit forming dense thickets dark red when right in fence rows, roadsicles "limon de chung" frents and all uncultivated places. eaten when night by dildren green fruits used for glue. (a serious pest, making walking very difficult of the trails). vine climbing in 08 abrus spresatorius emmon in thickets small trees, seeds scarlet with black end "kolales aniti (Devilsbeads) used for making necklaces. blowers frale-yellow. 09 Physalis lancealata
weed incultivated ground fruit edible, used medicinally. france fleshy galak" 10 Cyclophorus epiphytic on tree trunks (one of several plants of this name) leaves and rook briled, water taken for stomach ache. green roots plattened 11 Taeniophyllum epiphylic on tree-trunk against tree trunk "gamsan" (octopus grass) used to care pumples. tree 5 m. tall, flowers white, fruit pink, sweet "manzanita" servies eaten. common calabura

quam 149 148 # 1944 Marianas 1. small tree 4 m. talp. 25413 Leucaena glavea maken hotses hair ball out beans used but making mechanic and other maments. very abundant in Thickete everywhere used as fodder but leaves up to 1 m. a more 14 Fry John Xanthosoma high, blinish green, cultivated and spontaneous with a strong distinct in cultivated ground angle between petiole and blade, yautia" tuber used for food, young leaves also, as green regetable older leaves briled with coesnut milh and fed to prigs. (generally believed to come from Hawaii) chaquan fai (several grasses called this) common in open places used ber fordder tree 8 m stall, spreading, 16 fithecellobium dulce very ammon rounded, dense, flower white nave. "kamachili" fruit edible; cows and grats eat leaves; bark used for tanning, word used for house foundations; sometimes planted for hedges. shrub 4 m, tall, flower 17 Cestrum moctumum greenish white, very cultivated around church fragrant at night.

quan 151 1946 Marianas Do. 150 herb 1.5' m. tall, flowers yellow "amo tomaga carabao" 25418 Cassia occidentalis weed everywhere blue violet "dadalagesto 19 Stackytarpheta warm 1.) Vahl weed in distribed ground "seobilla anako" (because vo Lida acuta for browns. "mumuting umai" weed in open places 21 acalypha indica L. uneamon in open places flowers yellow "saigen" foolder only 22 Lynedrella nodiflora common in disturbed ground July 4 - hill between Merizo and Umatag gas south end of quam eroded volcanic soil twining vine with gray 23 Cantharospermum scarabaeoides (L.) Baill. leaves yellow flowers; very common "kakagustes aniti" (devils peanuts" leaves used medicinally. strongly aromatic. 24 Hyptis suaveoleus (L.) Pri. "mumutung" not used. when it grows vigorously it is thought to indicale feetel & erect, lastiferous, flowers 23 Euphorbia white "golondrina"

quam 153 1944 Marianas Is. 152 many decumbent to 25426 Phyllanthus ascending stems from emmon a root crown. July 6 - Comarianas, hills north of Mt. lenjo and mt. aluton, guan weedy disturbed places. largely thickets of Lewisena vine, flowers dell 27 I pomorea triloba L. rose purple. climbing in Whichets small tree 2 m. tall, 28 Bixa orellana Therb 1 m. tall. thicket 29 Walthering indica L. flowers yellow. armatic, flowers white. edge of thicket 30 agenatum congroides L 'edge of weedy thicket. suffruterent herbs 1.5 m. 31 Indigofera suffruticoa L. tall, flowers salmon pink. thickets flowers yellow, pods 32 Crotalaria guinquefolia L flowers yellow. weedy roadside 33 terseis Blumca simuata open weedy hilltop prostrate, flowers on bare ground. 34 Desmodium triflorum creeping on bare sports and among weeks, commen. slant grayish; blowers 39 Heliotropium oraklopulium common in disturbed places

mesophytic than that on Rota. It lach the Hernandia and has more species of non-strand plants. This vegetation is probably what logically may be expected when taybrical strand vegetation is elevated over a period of many years. The conditions tend to become less and less halofphytic as the formale rain-water washer the salt and ever alkali out. The continual inflers of propagreles that always taken place produces more and more effect - more and more invaders find conditions to their liking and become established. a succession taker place, enormously slow, toward a mesophytic condition. In such places cer differ, eliff tops, frest is not so well supported and

quan other more primeer species such as Phyllanthus mariannense, Hedyotis Goetidus, Bilsia, etc. are found. On the coastal plain, which is usually coal sand or slightly raised dissected coral transtranguer or low muddy swamp land, true strand conditions prevail. The usual fringe of Leaevola, Messersch-midia, Colubrina Casuarina, Hornoea per-caprae, mis, Wedelia, Carrytha, ct. lines the sandy portions the welly parts have Pemphys aciduly very abundant with 1 Leaevola, callicarpa, Hedyotis albido puntala Finhistylis cymosa, etc. Out in the rect flats in shallow salt water are Two see granes, 25322, 25323 very abundantly.

The low muddy areas have Bruguiera and Rhizophna, with Thesperia, a construction the edger. The land back of the beach supports a frest of Hobissus titiacens, Therperia, Hernandia, Intria bijuga, guettarda, Pithecellobium dula, Clerdendon, Caruaina, Cows musifera, Morinda c: tripolia, Vandanus tectorius, Clerodendon, etc. this tas in most places, especially where wet, been disturbed as much that it is hard to tell if any is really primary trand frest. This whole evental shelf seems to be a few feet of above sea level. and is being gradually end down, as evidenced by the remnants of ranged neef left here and there. The entling may well trave been

reduced now to that by solution and subalial erosin because of the development of a fringing Much of the subsistance agriculture of gram is on this cotastal shelf. Commuts are abundant, as are breadfruit, yams, Mangoes grow subspintaneously. Hogs and eattle and elichers are so apparently common normally, though rare In all distribut areas, of course, weeds are enously commer. The most prominent is Lemana glama which forms thickets everywhere and tends to develop ists a thick small firest. Tripliania trifolia runs et a close second, forming impenetrable tangler everywhere.

fram Marianas Is. 1946 One of the most distinctive Carica papara has developed regetation types is that of a small-fruited variety the areas of volcanic exposure which has truly become It is essentially a grassland a ubiquitous weed, especially which seems to be an early on disturbed limestone areas. successional stude. The deminant plant is miscanthus Japonius, which forms dense braker over much of the area. This is frequently burned off and erosion sears are a characteristic feature, in various stages of necovery. On these, and on ridges, rock outeroffs, and other places where the sword grass is either absent or sparse, there is a flora similar to that of the volcanie regions of Rota and failsan, but much riche in species. Several species of Fimbristylis, Helerofrogon, several other grasses, gleichenia linearis, Lyesfordium commen I tais sp. Blechnum sp. Lygodium scandens,

Wlithi 167 166 1944 Caroline 25. soil sandy, even as ich fuly 10 Fasserair Islet around village in evenut plantation onter beach prostrute forming patelres "pengapem" flower, purpliel, 25439 Centella asiatica (L.) Unh. 5 40 Vernonia cinerca (t.) Less. emmon spreading tree hulifini 3 41 Hibiseus Tiliaceus L. file of inner back Thickete used for making "grass" strute and worken was - arounds for women. tree 15 m. tall, buds 42 Engenia small grove and fruite few; tout Guit fleshy nather thite, seed free from pulp "haravath fruit eaten. 43 Prema cognitia R.Br. scattered small tree, flowers pale green, anomatec. very anomatic, for bracts . 44 Ocimum basilicum L. around houses and calyer pringlish evolla white "waring 45 Tacca leontopation (L)O.K.
commen everywhere ered acqueent heil, "mogmog" some say not.

Withi

July 11 - Mog mog Islet

gravelly substhatum

with a line of sattlet

broken coral at outer beach

flat, planted & coronut,

except large marsh

in center planted to

cyrtosperma, This

fillefwill much.

25419 Ochrosia opportifolia (Lam.) K. Schum.

1946

71 Melochia

island.

174

Caroline 1s.

, 69 allopphyllus nave , 70 guettarda speciova L. common

occasional on outer beach on flat that apparently has been cleared.

72 Vigna maring (burm.) Men.

common box on of sen

flats near channel beach

73 Pipturus argenteus

common on flats near

shannel and ruter books

seedling over whole

slender tree 4 m. tall,

flowers white; fruits

green, old ones priched

up under tree 4 m. tall,

flowers white.

flowers white, pleshy.

fruit white, pleshy.

flowers print, much

clarker in center.

fait of flowers graining

Caroline In. 184 1944 Ulithi stem flesly, reddish;
does not sting.
enlms spreading - ascending 25517 Fleurya underater gand. ummen 5 18 Parpalum common locally near march heads green 19 Eypens brenfoling I occasional ponds erect, stensively is Colypodium serloperdium emmen, especially creeping. on bases of trees plant flesling pruplish green. 21 Portulacy oberacea L. common maden distribus placer prostrate, root ealen "camole" spreading tufts 22 pomora batatas L. cultivated extensively 5 27 Enagrostis amabilis (L.) Weta: common to prostrate, fleshy 2 24 Vilea mierophylla (1) Lebm. common locally prostrate; flowers white, corolla Tube swollen 5 25 Hedystis-biflora abundant inopen places and under evenuts flowers white 4 26 Ocemen sanctum L doregard flowers yellow 5 * 19a Jussiaea suffentionea Mig.

1	188	Plants collected by Kulf. Pelzer			on Trule May- august, 1946189
	- 1	Ixora cacallinensis	2	16	Glockidion
		"acheu"	-		"amoses" Wona (Moen)
4	2	Polyscias grandifolia volla.	2	17	gleichema linearis
		"achenecha" Wona (Moen)			"onecha"
2	3	Crotalaria mucronata Desv.	2	1-3	
		afalafalu			ano" Wona (Moen)
7		Glochidion.	,	19	acrostichum aueum
		"afor "	4		"opössen" (Ton)
7	5	Cureuma	1	20	Acrostichum aureum
		"afan"	-		"apasou"
1	6	Parinarium glaberrimum	2	21	Eugenia
2		"ais"			"a"pe /" To/
. 7	7	Parinarium glaberrimum	4	72	
10 -		"eis" (ais) Tol			"apo" Wona (Noen), safein anu", safeian sa"t"
2	3	Hernandia ovigera	1	23	
		"akurang"	1-		"apuap"
7	9	Cordia subcordata	5	24	
		"alau" Toloas (Dublon)			"apuag"
	10	Cordia subcordata	1	25	Clevo dendron enerme
1116		"ano" Tol "alou" Moen	À		"apuech", "u/o"
. ,	11	Messer schmidia argentea		26	Ficus tinctoria var. neo-eludarum (summ.) Fosh.
No.		"a ma lose + " Wona (Moen)			"ouol", "apuilis" fruits are eaten
1	12	Nephro legis	3	27	Cassia alata
		"omare" (Vona (Moon)	1		"arakak" Tou (Tol)
2		Messerschmidia augentea	1	23	Terminolia catappa
18	-	"amaloset"			"0505"
11/1	14	Cyperus javanicus	1	29	Miscanthus floridulus
16		"a'ma'a" " emele"			"asat"
3	15		-	30	Randia Carolinensis (Val.) "Asepot" Tol, asepar Moen
N. Contraction		Veghrolegis "am a"re"	-		"Asepot" Tol, asepar "sen

190	191
100	
, 31 Wedelia biflora	4 44 Pemphis acidula
" a fux f"	engi Wona (Moen)
, 32 Wodelia biflora	1 47 Piper
"atuat"	"Ennes"
3 33 Jassiaea suffrinticosa L.	3 49 Piper
"aunenipain" Wona (Moen)	"Ennes" Wina (Moen)
34 Eroecaria agallocha	1 49 Dioscorea
"ansus" mangrove	"ep" Wona (Moen)
75 Ficus finctoria van. neo-elndarum (summ.) Forb.	
"macek" Moen, "awo'l" Tou (Tal)	Cleck on there notice
3 36 Psychotria carolineusis	1 51 £ names
chemei chemei	4 53
37 Psychofria carolineasis (val.) Fosb.	11 50 2 60
'Alamel " Lina (Mon)	" 77
Chemei Wona (Moen) 38 Solanam	1 53 / 1/2
	a divinuale
"chi" yellow fruit apples 29 Polypo dium scolopendria	2 54 Duglicating rumble
"Chichi"	62-
2 40 Colocasia esculenta	one is regentered in the other en)
"chi mök" Tou (Tol)	the freeres en)
2 41 Colocasia esculenta	2 56,
"Chimok" ploen	(Moen)
6 42 Randia carolinensis	t 57
"chomis" - Toloas (Dublon)	
	3 58
"pirenges" Tol, "one put "Moen; small white flowers	
41.079	3 59 Jussiaea suffruiticosa
1 49 Ipomora digitata	"likainenpal" Tou (Tol)
"emechimech" Tol; nechimech Moon	
r 45 Glockickion "emoses"	"liko ioio" Tou (Tol) Roxb.

Cleck on There retire names. 112 Duglicating number one if regestered in the # book the other is cyperus

				.=>:		
1	15	92	62 a Cypelus			193
			"manon" Operat Wing			
1	1 =		Hibiscus moschatus wina (moen)	1	7,6	Cassia
			"liko no kon" - Tou (To!) Lopeanen amara			"Nikamaur"
	1	62	30 tila mea amara	١	77	Scleria
1 =	-		"Maras"- Moon "pwanu"-Tol	7		"Mika unaun" - Wona (Moen) Nika safasap" Toutsel)
3	97		Ficus tinctoria vac. neo-chridanim (Sum.) Forb.		78	Mounda citrifolia
1/4			Mauk - "Wona" (Moen)		79	Nopur"
7	V		Davallia solida			Morrida cifrifolia
			"mechenkafu"	ic	51	"Nupur" - Moen, "Arin". Tou (Tol) Cynometra.
	1	65	Ipomoca digitata.		-	"omo"
W			mechimoch (Ost) emechimech (west)	2	81	Cynometra
	1		"Mongif"			"omio" Toloas (Dublon)
	0		Pisonia grandis	2	82	Bruguiera conjugata
	-	6/	"mök" Wöna (Moen)			"ong"
14	2	68	Amorphophallus	L	83	
	-		"Moku fen"	1		"ougin"
	2	69	Euphobia hirta	1	84	Agera fum cony 201 des
101			"kichich" - Tol, Moneniap - Moen	-		"opo"nu pon " Toa (Tol)
	2	70	Kleinhouia hospita	3	85	Piper
HIL.			Monou Wona (Moca)	+-	C	"ötil" wona (Moen)
	1	71	Luffa cylindrica	1		Fagraea
			"mororof" Tol, Moen			"O-ur", sour Moen
MIN	5	72	Gueffordsa speciosa	1	87	"our", a-ur Tol, so-ur Moen
		-	"MOSOF" Wo"na (Moen)	1	66	
	1-		Allophylus		00	"par" Tou (Tol)
		74	"Ngo" Wona (Moen)	2	£5	Ery thrina
	1		Allophylus (TI)		0 1	"Par"
	0	75	Ngo"-Tou (Tol)	. 2	90	
W	1		"Nika chop a chop " Works (Moon); Grows on Mai, Kuz, Vakich			"pula pul", Penina) - Moen



